
UNRAMIFIED CORRESPONDENCES

by

Fedor Bogomolov and Yuri Tschinkel

ABSTRACT. — We study correspondences between algebraic curves defined over the separable closure of $\mathbb{Q}$ or $\mathbf{F}_p$.

Contents

Introduction	1
2. Main construction	3
3. The case of characteristic 0	6
4. Geometric constructions	8
References	12

Introduction

A class $\mathcal{C}(\overline{\mathbb{Q}})$ of complete algebraic curves over $\overline{\mathbb{Q}}$ will be called *dominating* if for every algebraic curve C' over $\overline{\mathbb{Q}}$ there exist a curve $\tilde{C} \in \mathcal{C}(\overline{\mathbb{Q}})$ and a birational surjective map $\tilde{C} \rightarrow C'$. A curve C will be called *universal* if the class $\mathcal{U}_C(\overline{\mathbb{Q}})$ of its unramified covers is dominating.

THEOREM 1.1 (Belyi). — *Every algebraic curve C defined over a number field admits a surjective map onto $\mathbb{P}^1$ which is unramified outside $(0, 1, \infty)$.*

In 1978 Manin pointed out that Belyi's theorem implies the following

PROPOSITION 1.2. — *The class $\mathcal{MU}(\overline{\mathbb{Q}})$ consisting of modular curves and their unramified covers is dominating.*

There are many other classes of curves with the same property, for example:

1. hyperelliptic curves and their unramified coverings;
2. the class $\mathcal{CU}(\overline{\mathbb{Q}}) := \cup_{n \in \mathbb{N}} \mathcal{C}_n(\overline{\mathbb{Q}})$, with $\mathcal{C}_n(\overline{\mathbb{Q}})$ consisting of curves with function field $\overline{\mathbb{Q}}(z, \sqrt[n]{z(1-z)})$ and their unramified coverings.
3. the class $\mathcal{CN}(\overline{\mathbb{Q}}) := \cup_{n \in \mathbb{N}} \mathcal{CN}_n(\overline{\mathbb{Q}})$ where $\mathcal{CN}_n(\overline{\mathbb{Q}})$ consists of all unramified covers of any curve C_n with the property that $C_n \rightarrow \mathbb{P}^1$ is ramified in $(0, 1, \infty)$ only and all local ramification indices of C_n over 0 are divisible by 3, over 1 divisible by 2 and over ∞ divisible by n . In particular, we could take C_n to be the modular curve $X(n)$.

Proof. — (Sketch) Let us consider the class of hyperelliptic curves and their unramified covers. Let C' be an arbitrary curve and $\sigma : C' \rightarrow \mathbb{P}^1$ a *generic* map, branched over the points $q_1, \dots, q_n$ (generic means that there is only one ramification point over each branch point and all local ramification indices are equal to 2). Denote by C a hyperelliptic curve whose ramification contains $q_1, \dots, q_n$. Then $\tilde{C} := C \times_{\mathbb{P}^1} C'$ is an unramified cover of C which surjects onto C' . For the classes $\mathcal{CU}(\overline{\mathbb{Q}})$ and $\mathcal{CN}(\overline{\mathbb{Q}})$ we use Belyi's theorem. $\square$

QUESTION 1.3. — Does there exist a universal algebraic curve C (over $\overline{\mathbb{Q}}$)?

QUESTION 1.4. — Does there exist a number $n \in \mathbb{N}$ such that every curve defined over $\overline{\mathbb{Q}}$ admits a surjective map onto $\mathbb{P}^1$ with ramification over $(0, 1, \infty)$ such that all local ramification indices are $\leq n$?

QUESTION 1.5. — Is every curve C (over $\overline{\mathbb{Q}}$) of genus $g(C) \geq 2$ universal?

REMARK 1.6. — It is clear that an affirmative answer to Question 1.4 implies a (constructive) affirmative answer to Question 1.3.

In this note we answer these questions in a simple model situation: instead of $\overline{\mathbb{Q}}$ we consider the (separable) closure $\overline{F}_p$ of the finite field $\mathbf{F}_p$.

THEOREM 1.7. — *Let $p \geq 5$ be a prime and C a hyperelliptic curve over $\overline{\mathbf{F}}_p$ of genus $g(C) \geq 2$. Then C is universal: for any projective curve C' there exist a finite étale cover $\tilde{C} \rightarrow C$ and a surjective regular map $\tau : \tilde{C} \rightarrow C'$.*

In Section 4 we prove the following geometric fact (over arbitrary algebraically closed fields of characteristic $\neq 2, 3$):

PROPOSITION 1.8. — *Every hyperelliptic curve C has a finite étale cover $\tilde{C}$ which surjects onto the genus 2 curve C_0 given by $\sqrt[6]{z(1-z)}$. In particular, if C_0 is universal then every hyperelliptic curve of genus ≥ 2 is universal.*

REMARK 1.9. — Applying the Chevalley-Weil theorem we conclude that the Mordell conjecture (Faltings' theorem) for C_0 implies the Mordell conjecture for every hyperelliptic curve of genus ≥ 2 .

The fact that there is some interaction between the arithmetic of different curves has been noted previously. Moret-Bailly and Szpiro showed (see [6], [5]) that the proof of an *effective* Mordell conjecture for *one* (hyperbolic) curve (for example, C_0) implies the ABC-conjecture, which in turn implies an effective Mordell conjecture for *all* (hyperbolic) curves (Elkies [4]). Here *effective* means an explicit bound on the height of a K -rational point on the curve for all number fields K . Here again, Belyi's theorem is used in an essential way.

Acknowledgments. We have benefited from conversations with B. Hassett and A. Chambert-Loir. The first author was partially supported by the NSF. The second author was partially supported by the NSF and the Clay foundation.

2. Main construction

NOTATIONS 2.1. — Let $\tau : C \rightarrow C'$ be a surjective map of algebraic curves. We denote by $\text{Ram}(\tau) \subset C$ the ramification locus of τ and by

$\text{Bran}(\tau) = \tau(\text{Ram}(C)) \subset C'$ the branch locus of τ . For a point $q \in C$ we denote by $e_q(\tau)$ the local ramification index at q . We denote by

$$e(\tau) := \max_{q \in C} e_q(\tau)$$

the maximum local ramification index of τ . We say that τ has *simple* ramification if $e(\tau) \leq 2$ and that τ is *generic* if in addition there is only one ramification point over each branch point.

REMARK 2.2. — Every curve admits a generic map onto $\mathbb{P}^1$, at least after a separable extension of the ground field.

Let $p \geq 5$ be a prime number. In this section we work over a separable closure $\overline{\mathbf{F}}_p$ of the finite field $\mathbf{F}_p$. First we show that there exists at least one curve satisfying the conclusion of Theorem 1.7.

Let $\pi_0 : E_0 \rightarrow \mathbb{P}^1$ be the elliptic curve given by

$$\sqrt[3]{z(z-1)}.$$

Let $\sigma_0 : C_0 \rightarrow \mathbb{P}^1$ be the genus 2 curve given by

$$\sqrt[6]{z(z-1)},$$

and $\iota_0 : C_0 \rightarrow E_0$ the corresponding 2-cover. Clearly, ι_0 has simple ramifications over the preimages of 0, 1. Let C be an arbitrary curve. Choosing a generic function on C we get a generic covering $\sigma : C \rightarrow \mathbb{P}^1$ (such covering is defined over $\overline{\mathbf{F}}_p$). Assume further that $\text{Bran}(\sigma) \subset \mathbb{P}^1$ does not contain $(0, 1, \infty)$.

Consider the diagram

$$\begin{array}{ccccc} C & \longleftarrow & C_1 & \longleftarrow & C_2 \\ \sigma \downarrow & & \downarrow & & \downarrow \\ \mathbb{P}^1 & \longleftarrow & E_0 & & \\ & & \varphi \downarrow & & \\ & & E_0 & \longleftarrow & C_0 \end{array}$$

Here $C_1 = C \times_{\mathbb{P}^1} E_0$ (it is irreducible since $E_0 \rightarrow \mathbb{P}^1$ is a 2-cover). Then $C_1 \rightarrow E_0$ has simple ramification over a finite number of points

in E_0 . Recall that E_0 has a group scheme structure, and *all* $\overline{\mathbf{F}}_p$ -points of E_0 are torsion points. This implies that there exists an étale map $E_0 \rightarrow E_0$ such that all ramification points of C_1 over E_0 are mapped to 0. More precisely, any finite set of $\overline{\mathbf{F}}_p$ -points of E_0 is contained in the group subscheme $E_0^{et}[n] \subset E_0$ - the maximal étale subgroup of the multiplication by n -kernel $E_0[n]$ (for some $n \in \mathbb{N}$). For every positive integer n there exists a positive multiple m of n and an étale map $E_0 \rightarrow E_0$ with kernel $E_0^{et}[m]$.

Taking the composition of $C_1 \rightarrow E_0$ with the multiplication by a suitable m , we get a (possibly new) surjective regular map $C_1 \rightarrow E_0$ which is ramified only over the zero point in E_0 and has the property that all the local ramification indices are at most 2. Using this map let us define $C_2 := C_0 \times_{E_0} C_1$. Consequently, any component of C_2 surjects onto C_1 and is an étale covering of C_0 (ramification cancels ramification). This component satisfies the conclusion of Theorem 1.7.

LEMMA 2.3. — *Let C be any smooth complete algebraic curve and E any curve of genus 1. There exists a curve C_1 which surjects onto C and E such that the ramification of the map $C_1 \rightarrow E$ lies entirely over a single point of E and its local ramification indices are all equal to 2.*

Proof. — Consider a generic map $\sigma : C \rightarrow \mathbb{P}^1$ with $e(\sigma) \leq 2$. Choose a double cover $\pi : E \rightarrow \mathbb{P}^1$ such that the branch loci $\text{Bran}(\sigma)$ and $\text{Bran}(\pi)$ on $\mathbb{P}^1$ are disjoint. Then the product $C_1 := C \times_{\mathbb{P}^1} E$ is an irreducible curve which is a double cover of C . The curve admits a surjective map $\iota_1 : C_1 \rightarrow E$ with $e(\iota_1) \leq 2$. Similarly to the previous construction we can find an unramified cover $\varphi : E \rightarrow E$ such that the composition $\varphi \circ \iota_1 : C_1 \rightarrow E$ is ramified only over one point in E and the local ramification indices are still equal to 2. $\square$

COROLLARY 2.4. — *Assume that some unramified covering $\tilde{C}$ of C surjects onto an elliptic curve E . Assume further that there exists a point q on E such that all local ramification indices of the map $\tilde{C} \rightarrow E$ over q are divisible by 2. Then C is universal.*

Proof. — It is sufficient to take the product of $\tilde{C} \times_E C_1$. Any irreducible component of the resulting curve will be an unramified covering of $\tilde{C}$ (and hence C) and will admit a surjective map onto C_1 and C . $\square$

COROLLARY 2.5 (Theorem 1.7). — *Every hyperelliptic curve C over $\overline{\mathbf{F}}_p$ (with $p \geq 5$) of genus ≥ 2 is universal.*

Proof. — Consider the standard projection $\sigma : C \rightarrow \mathbb{P}^1$ (of degree 2). Its branch locus $\text{Bran}(\sigma)$ consists of $2g + 2$ points. Let $\pi : E \rightarrow \mathbb{P}^1$ be a double cover such that $\text{Bran}(\pi)$ is contained in $\text{Bran}(\sigma)$. Then the product $\tilde{C} = C \times_{\mathbb{P}^1} E$ is an unramified double cover of C . Moreover, $\tilde{C}$ is a double cover of E with ramification at most over the preimages in E of the points in $\text{Bran}(\sigma) \setminus \text{Bran}(\pi)$. We now apply Corollary 2.4. $\square$

In *finite* characteristic, there are many other (classes of) universal curves. For example, cyclic coverings with ramification in 3 points, hyperbolic modular curves, etc. Thus it seems plausible to formulate the following

CONJECTURE 2.6. — Any smooth complete curve C of genus $g(C) \geq 2$ defined over $\overline{\mathbf{F}}_p$ (for $p \geq 2$) is universal.

3. The case of characteristic 0

In this section we work over $\overline{\mathbb{Q}}$. We show that the method outlined in Section 2 can be employed in characteristic zero to produce natural infinite sets of algebraic points on $\mathbb{P}^1$ which occur as ramification points of surjective maps from $\mathbb{P}_2^1$ to $\mathbb{P}_1^1$ branched over $(0, 1, \infty) \in \mathbb{P}_1^1$ only and having an *a priori* bound on the ramification index (here $\mathbb{P}_1^1$ and $\mathbb{P}_2^1$ are two different copies of the projective line $\mathbb{P}^1$).

Notice that, in principle, it is easy to produce *some* sets of points (of any finite cardinality) with this property: Take an $n \geq 6$ and any triangulation of $\mathbb{P}_2^1$ with vertices of index $\leq n$. A barycentric subdivision of each such triangulation defines a function from $\mathbb{P}_2^1$ to $\mathbb{P}_1^1$ with local ramification indices $\leq 2n$ (for more details see [3]). Therefore, any curve with

bounded ramification over this set of vertices will have bounded ramification over $\mathbb{P}_1^1$. However, we have no explicit control over the coordinates of the ramification points on $\mathbb{P}_2^1$.

An (obvious) analogous way to control ramification indices is to consider the following diagram

$$\begin{array}{ccc} E & \xrightarrow{\pi} & \mathbb{P}_2^1 \\ \phi_n \downarrow & & \downarrow \varphi_{n,E} \\ E & \xrightarrow{\pi} & \mathbb{P}_1^1 \end{array}$$

where the map ϕ_n is the quotient by the subscheme of n -torsion points and the maps $E \rightarrow \mathbb{P}^1$ are the standard double covers, ramified over $(0, 1, \infty, \lambda)$. Clearly, all the ramification points of $\varphi_{n,E}$ (in $\mathbb{P}_2^1$) are over $0, 1, \infty$ and λ (in $\mathbb{P}_1^1$) and $e(\varphi_{n,E}) = 2$. Belyi's theorem gives a map $\beta : \mathbb{P}_1^1 \rightarrow \mathbb{P}_0^1$, which ramifies only over the points $(0, 1, \infty) \in \mathbb{P}_0^1$, maps $\{0, 1, \infty, \lambda\} \subset \mathbb{P}_1^1$ into $\{0, 1, \infty\} \subset \mathbb{P}_0^1$ and has local ramification indices $\leq n$. Moreover, it provides an explicit bound on $\deg(\beta)$ and, consequently, on $e(\beta)$ (in terms of the absolute height of λ). Let $\beta_\lambda : \mathbb{P}_1^1 \rightarrow \mathbb{P}_0^1$ be a map such that

$$e(\beta_\lambda) = \inf_{\beta} \{e_{\beta}\}$$

over the set of all maps as above. Then the map $\beta_\lambda \circ \varphi_{n,E} : \mathbb{P}_2^1 \rightarrow \mathbb{P}_0^1$ ramifies over three points only and has index $e(\beta_\lambda \circ \varphi_{n,E}) \leq 2n$. Let

$$R_E := \pi(E(\overline{\mathbb{Q}})_{\text{tors}}) \subset \mathbb{P}_2^1(\overline{\mathbb{Q}})$$

be the image of the torsion points of E . Let $\sigma : C \rightarrow \mathbb{P}_2^1$ be any map ramified only in a subset of R_E . Let $\pi := \beta_\lambda \circ \varphi_{n,E} \circ \sigma$. Then

$$e(\pi) \leq 2e(\sigma) \cdot e(\beta_\lambda).$$

A natural application of the construction in Section 2 is as follows:

EXAMPLE 3.1. — Let $\pi : E \rightarrow \mathbb{P}^1$ be a triple cover with $\text{Bran}(\pi) = \{0, 1, \infty\}$ (E is a CM elliptic curve with j -invariant 0). Consider the following diagram

$$\begin{array}{ccc}
E & \xrightarrow{\pi} & \mathbb{P}_2^1 \\
\phi_n \downarrow & & \downarrow \varphi_{n,E} \\
C_0 \longrightarrow E & \xrightarrow{\pi} & \mathbb{P}_1^1,
\end{array}$$

where C_0 is a curve of genus $g(C_0) = 2$ given by $\sqrt[6]{z(z-1)}$, ϕ_n is the quotient map by the subscheme of torsion points of order n , and $\varphi_{n,E}$ the corresponding map from $\mathbb{P}_2^1$ to $\mathbb{P}_1^1$ ramified only over $(0, 1, \infty)$. Let $\mathcal{X}_g = \{X\}$ be the subset of curves of genus g admitting a map $\sigma_X : X \rightarrow \mathbb{P}_2^1$ such that

- $e(\sigma_X) = 2$;
- $\text{Bran}(\sigma_X) \subseteq \pi(E(\overline{\mathbb{Q}})_{\text{tors}})$.

Then, for any $X \in \mathcal{X}_g$ the map

$$\varphi_{n,E} \circ \sigma_X : X \rightarrow \mathbb{P}_1^1$$

has index $e(\varphi_{n,E} \circ \sigma_X) \leq 6$ and there exists an unramified cover $\tilde{C} \rightarrow C_0$ surjecting onto X . Moreover, $\mathcal{X}_g$ is *dense* (in real and p -adic topologies) in the natural Hurwitz scheme $\mathcal{H}_g$ parametrizing curves of genus g .

The set of curves dominated by unramified covers of C_0 is much larger than $\mathcal{X}_g$. Indeed, consider any 4-tuple of points in

$$\pi(E(\overline{\mathbb{Q}})_{\text{tors}}) \subseteq \mathbb{P}_2^1$$

and an elliptic curve E' obtained as a double cover of $\mathbb{P}_2^1$ ramified in those 4 points. Then E' is also dominated by unramified covers of C_0 and we can iterate the above construction for E' .

4. Geometric constructions

Let (E, q_0) be an elliptic curve, q_1 a torsion point of order two on E and $\pi : E \rightarrow \mathbb{P}^1$ the quotient with respect to the involution induced by q_1 . Let n be an odd positive integer and $\varphi_{n,E} : \mathbb{P}_2^1 \rightarrow \mathbb{P}_1^1$ the map induced by

$$\begin{array}{ccc}
E & \xrightarrow{\pi} & \mathbb{P}_2^1 \\
\phi_n \downarrow & & \downarrow \varphi_{n,E} \\
E & \xrightarrow{\pi} & \mathbb{P}_1^1.
\end{array}$$

Any quadruple $r = \{r_1, \dots, r_4\}$ of four distinct points in $\varphi_{n,E}^{-1}(\pi(q_0))$ defines a genus 1 curve E_r (the double cover of $\mathbb{P}^1$ ramified in these four points).

PROPOSITION 4.1. — *Let $\iota : C \rightarrow E$ be any finite cover such that all local ramification indices over q_0 are even. Then there exists an unramified cover $\tau_r : C_r \rightarrow C$ which dominates E_r and has only even local ramification indices over some point in E_r .*

Proof. — Assume that $n \geq 3$ and consider the following diagram

$$\begin{array}{ccccc}
C & \xleftarrow{\tau_2} & C_2 & \xleftarrow{\tau_r} & C_r \\
\downarrow \iota & & \downarrow \iota_2 & & \downarrow \iota_r \\
E & \xleftarrow{\varphi_n} & E & & E_r \\
\downarrow \pi & & \downarrow \pi & & \downarrow \pi_r \\
\mathbb{P}_1^1 & \xleftarrow{\phi_{n,E}} & \mathbb{P}_2^1 & & \mathbb{P}_2^1,
\end{array}$$

where E_r is a double cover of $\mathbb{P}_2^1$ ramified in any quadruple of points in the preimage $\phi_{n,E}^{-1}(\pi(q_0))$ and C_r is any irreducible component of $C_2 \times_{\mathbb{P}_2^1} E_r$. Any point $q_r \in E_r$ such that $q_r \notin \text{Ram}(\pi_r)$ (that is, its image in $\mathbb{P}_2^1$ is distinct from $r_1, \dots, r_4$) has the claimed property. $\square$

REMARK 4.2. — Iterating this procedure (and adding isogenies) we obtain many elliptic curves E' which are dominated by curves having an unramified cover onto E . It would be interesting to know if for any two elliptic curves over $\overline{\mathbb{Q}}$ there exists a cycle connecting them (at least modulo isogenies). We will now show that *any* elliptic curve can be connected in this way to E_0 .

Let $E_0 \subset \mathbb{P}^2 = \{(x : y : z)\}$ be the elliptic curve

$$x^3 + y^3 + z^3 = 0,$$

and

$$E_0[3] = \mathsf{T} := \left\{ \begin{array}{lll} (1 : 0 : 1), & (1 : 0 : -\zeta), & (1 : 0 : -\zeta^2), \\ (0 : 1 : 1), & (0 : 1 : -\zeta), & (0 : 1 : -\zeta^2), \\ (1 : 1 : 0), & (1 : -\zeta : 0), & (1 : -\zeta^2 : 0) \end{array} \right\}$$

its set of 3-torsion points (where ζ is a primitive cubic root of 1). Denote by $\mathcal{E}_\lambda = \{E_\lambda\}$ the family of elliptic curves on $\mathbb{P}^2$ passing through T given by

$$E_\lambda : x^3 + y^3 + z^3 + \lambda xyz = 0.$$

It is easy to see that for each λ the set $E_\lambda[3]$ of 3-torsion points of E_λ is precisely T .

$$\begin{array}{ccc} \pi & : & \mathbb{P}^2 \quad \rightarrow \quad \mathbb{P}^1 \\ & & (x : y : z) \mapsto (x + z : y) \end{array}$$

the projection respecting the involution $x \rightarrow z$ on $\mathbb{P}^2$. Denote by π_λ the restriction of π to E_λ . Clearly, π_λ exhibits each E_λ as a double cover of $\mathbb{P}^1$ and π_λ has only simple double points for all λ . Moreover,

$$\pi(\mathsf{T}) = \{(0 : 1), (1 : -\zeta), (1 : -\zeta^2), (1 : -1), (1 : 0)\}$$

and for all λ there exists a (non-empty) set $S_\lambda \subset \text{Bran}(\pi_\lambda) \subset \mathbb{P}^1$ such that $\pi_\lambda^{-1}(S_\lambda) \subset \mathsf{T}$. Let $\pi'_0 : E'_0 \rightarrow \mathbb{P}^1$ be a double cover ramified in 4 points in $\pi(\mathsf{T})$.

LEMMA 4.3. — *Let $\iota : C \rightarrow E_\lambda$ be a double cover such that over at least one point in $\text{Bran}(\iota)$ the local ramification indices are even. Then there exists an unramified cover $\tilde{C} \rightarrow C$ and a surjective morphism $\tilde{\iota} : \tilde{C} \rightarrow E'_0$ such that over at least one point in $\text{Bran}(\tilde{\iota}) \subset E'_0$ all local ramification indices of $\tilde{\iota}$ are even.*

Proof. — Consider the diagram

$$\begin{array}{ccc} E_\lambda & \xleftarrow{\iota} & C_1 \\ \varphi_3 \downarrow & & \downarrow \\ E_\lambda & \xleftarrow{\quad} & C \\ \pi_\lambda \downarrow & & \\ & & \mathbb{P}^1 \end{array}$$

Then $C_1 \rightarrow \mathbb{P}^1$ has even local ramification indices over all points in $\pi(\mathbb{T})$. It follows that

$$\tilde{C} := C_1 \times_{\mathbb{P}^1} E'_0 \rightarrow E'_0$$

has even local ramification indices over the preimages of the fifth point in $\pi(\mathbb{T})$, as claimed. $\square$

NOTATIONS 4.4. — Let $\mathcal{C}$ be the class of curves such that there exists an elliptic curve E , a surjective map $\iota : C \rightarrow E$ and a point $q \in \text{Bran}(\iota)$ such that all local ramification indices at points in $\iota^{-1}(q)$ are even.

EXAMPLE 4.5. — Any hyperelliptic curve of genus ≥ 2 belongs to $\mathcal{C}$. More generally, $\mathcal{C}$ contains any curve C admitting a map $C \rightarrow \mathbb{P}^1$ with even local ramification indices over at least 5 points in $\mathbb{P}^1$.

PROPOSITION 4.6. — *For any $C \in \mathcal{C}$ there exists an unramified cover $\tilde{C} \rightarrow C$ surjecting onto C_0 (with $C_0 \rightarrow \mathbb{P}^1$ given by $\sqrt[6]{z(1-z)}$).*

Proof. — Consider $C_1 = C \in \mathcal{C}$ with $\iota_1 : C_1 \rightarrow E = E_\lambda$ as in 4.4. Define C_2 as an irreducible component of $C_1 \times_E E$:

$$\begin{array}{ccc} C_1 & \xleftarrow{\tau_2} & C_2 \\ \iota_1 \downarrow & & \downarrow \iota_2 \\ E & \xleftarrow{\varphi_3} & E \\ & & \downarrow \pi_\lambda \\ & & \mathbb{P}^1 \end{array}$$

Define $C_3 := C_2 \times_{\mathbb{P}^1} E_0$ by the diagram

$$\begin{array}{ccc} C_2 & \xleftarrow{\tau_3} & C_3 \\ \sigma_2 \downarrow & & \downarrow \iota_3 \\ \mathbb{P}^1 & \xleftarrow{\pi_0} & E_0. \end{array}$$

Observe that for $q \in \text{Bran}(\pi_0)$ the local ramification indices in the preimage $(\iota_2 \circ \pi_\lambda)^{-1}(q)$ are all even. It follows that the map $\tau_3 : C_3 \rightarrow C_2$

is *unramified* and that $\iota_3 : C_3 \rightarrow E_0$ has even local ramification indices over (the preimage of) $q_5 \in \{\pi(\mathbf{T}) \setminus \text{Bran}(\pi_0)\}$ (the 5th point). Define C_4 as an irreducible component of $C_3 \times_{E_0} E_0$ in the diagram

$$\begin{array}{ccc} C_3 & \xleftarrow{\tau_4} & C_4 \\ \iota_3 \downarrow & & \downarrow \iota_4 \\ E_0 & \xleftarrow{\varphi_3} & E_0. \end{array}$$

The map ι_4 is ramified over the preimages $(\pi_0 \circ \varphi_3)^{-1}(q_5)$, with even local ramification indices. Finally, $C_5 = C_4 \times_{E_0} C_0$ from the diagram

$$\begin{array}{ccc} C_4 & \xleftarrow{\tau_5} & C_5 \\ \iota_4 \downarrow & & \downarrow \\ E_0 & \xleftarrow{\iota_0} & C_0. \end{array}$$

has a dominant map onto C_0 and is unramified over C_4 (and consequently, C_1). $\square$

References

- [1] G. V. Belyi, *Galois extensions of a maximal cyclotomic field*, Izv. Akad. Nauk SSSR Ser. Mat. **43**, (1979), no. 2, 267–276, 479.
- [2] G. V. Belyi, *Another proof of the Three Points theorem*, Preprint MPI 1997-46 at <http://www.mpim-bonn.mpg.de>, (1997).
- [3] F. Bogomolov, D. Husemoller, *Geometric properties of curves defined over number fields*, Preprint MPI 2000-1 at <http://www.mpim-bonn.mpg.de>, (2000).
- [4] N. Elkies, *ABC implies Mordell*, Intern. Math. Res. Notices **7**, (1991), 99–109.
- [5] L. Moret-Bailly, *Hauteurs et classes de Chern sur les surfaces arithmétiques*, Astérisque **183**, (1990), 37–58.
- [6] L. Szpiro, *Discriminant et conducteur des courbes elliptiques*, Astérisque **183**, (1990), 7–18.